Contents

Preface vii

Conference information viii

Conference photograph xi

Chapter 1. Integer quantum Hall effect

The quantum Hall effect – an edge phenomenon?
K. von Klitzing 1

Nonequilibrium electron distribution and nonlocal resistance in a two-dimensional electron gas at high magnetic fields
S. Komiyama and H. Nii 7

Device-width dependence of plateau width in quantum Hall states
S. Kawaji, K. Hirakawa and M. Nagata 17

Influence of gate voltage on nonlocal resistance in GaAs/AlGaAs heterostructure at high magnetic fields
S. Takaoka, K. Tsukagoshi, K. Oto, K. Murase, Y. Takagaki, K. Gamo and S. Namba 21

Anderson transition in layered systems in quantizing magnetic fields
T. Ohtsuki, B. Kramer and Y. Ono 26

Quantized Hall effect in 3D periodic systems
M. Kohmoto, B.I. Halperin and Y.-S. Wu 30

Phase-coherence length in a two-dimensional electron gas at high magnetic fields
H. Hirai, S. Komiyama, K. Nakamura and F. Nihey 34

Chapter 2. Fractional quantum Hall effect and Wigner solid

Wigner solid versus incompressible Fermi liquid: phase diagram derived from time-resolved photoluminescence
I.V. Kukushkin, N.J. Pulsford, K. von Klitzing, R. Haug, K. Ploog and V.B. Timofeev 38

Fractional quantum Hall effect in multilayer electron systems
Y.W. Suen, J. Jo, M.B. Santos, L.W. Engel and M. Shayegan 43

Correlated 2D electrons in high magnetic fields
G. Martinez 51

High-field photoluminescence in GaAs single heterojunctions
A.G. Davies, S.A. Brown, R.B. Dunford, E.M. Goldys, R. Newbury, R.G. Clark, P.E. Simmonds, J.J. Harris and C.T. Foxon 56

Incompressible quantum liquid versus quasi-two-dimensional electron solid
G. Meissner 66

Scaling in the metal–insulator transition of the fractional quantum Hall effect
S. Koch, R.J. Haug, K. von Klitzing and K. Ploog 72

Observation of a spin-polarization instability in tilted magnetic fields
S. Koch, R.J. Haug, K. von Klitzing and M. Razeghi 76

Observation of the fractional quantum Hall effect in GaAs–(Ga,Al)As quantum well structures
C.V. Brown, C.J.G.M. Langerak, P.C. Main, L. Eaves, T.J. Foster, M. Henini, P.A.A. Teunissen and J.A.A.J. Perenboom 81

Fractional quantum Hall effect in narrow channels
D. Yoshioka 86

Electronic structure of a double fractional quantum Hall system of spin-1/2 electrons
T. Nakajima and H. Aoki 91

Observation of the transition to an insulating state consistent with a Wigner solid in a high-density 2D hole gas
P.J. Rodgers, C.J.G.M. Langerak, B.L. Gallagher, R.J. Barraclough, M. Henini, T.J. Foster, G. Hill, S.A.J. Wiegers and J.A.A.J. Perenboom 95

Chapter 3. Magneto-optics

Collective excitations in a quasi-3-dimensional electron system
H.D. Drew, X. Ying, K. Karraï, M. Shayegan and M. Santos 100

A magneto-optical study of coupled quantum wells in strained GaInSb/GaSb
S.L. Wong, R.J. Warburton, R.J. Nicholas, N.J. Mason and P.J. Walker 106

New features in magneto-optical spectra of $GaAs/Al_xGa_{1-x}As$ short-period superlattices observed under pulsed high magnetic fields up to 150 T
S. Sasaki and N. Miura 111

Many-body effects on magneto-luminescence due to acceptor states in semiconductor quantum wells
S. Katayama and T. Ando 115

The high magnetic field dependency of 2D exciton states in GaAs/AlAs multi-quantum-wells
T. Yasui, Y. Segawa, Y. Iimura, Y. Aoyagi, I. Mogi and G. Kido 119

Stimulated Landau emission by EMF excitation with application to spectroscopy of InSb at the quantum limit
T. Morimoto, M. Chiba, S. Ueda, G. Kido and M. Inoue 123

High magnetic field effects on electron–bound hole recombination in InSb
H. Kobori, T. Ohyama and E. Otsuka 128

Magneto-optical study on the excitonic spectrum of $(C_6H_{13}NH_3)_2PbI_4$
T. Kataoka, T. Kondo, R. Ito, S. Sasaki, K. Uchida and N. Miura 132

Zeeman effect on the zero-phonon line of the NV center in synthetic diamond
H. Hanzawa, H. Nishikori, Y. Nisida, S. Sato, T. Nakashima, S. Sasaki and N. Miura 137

Chapter 4. Cyclotron resonance

Optically detected cyclotron resonance in Ge, Si and ZnSe
T. Ohyama, T. Tomaru and E. Otsuka 141

Cyclotron resonance of mixed phases of the 2-D system in the ultra-quantum limit
G.M. Summers, M. Watts, R.A. Lewis, R.J. Nicholas, J.J. Harris and C.T. Foxon 149

Cyclotron resonance measurements of the hole mass in [0 0 1] and [1 1 1] $In_xGa_{1-x}Sb/GaSb$ quantum wells
R.J. Warburton, R.A. Lewis, R.J. Nicholas, N.J. Mason and P.J. Walker 154

Optically detected cyclotron resonance of GaAs quantum wells
J.G. Michels, R.J. Warburton, R.J. Nicholas, J.J. Harris and C.T. Foxon 159

Magneto-optical and far-infrared optically detected cyclotron resonance determination of the effective mass in $GaAs_{1-x}P_x$
P. Omling, C. Wetzel, A.L. Efros and B.K. Meyer 164

Ultra-high magnetic field cyclotron resonance of zero-gap InAs/GaSb superlattices
D.J. Barnes, R.J. Nicholas, N.J. Mason, P.J. Walker, R.J. Warburton and N. Miura 168

Anomalous temperature dependence of the effective mass in PbTe and PbSe
H. Yokoi, S. Takeyama, O. Portugall, N. Miura and G. Bauer 173

Observation of cyclotron resonance in low-mobility semiconductors using pulsed ultra-high magnetic fields
J. Kono, N. Miura, S. Takeyama, H. Yokoi, N. Fujimori, Y. Nishibayashi, T. Nakajima, K. Tsuji and M. Yamanaka 178

Chapter 5. Magneto-transport phenomena

Magneto-transport and magneto-optics in narrow-gap III–V compounds based on InSb
T. Le, W.T. Yuen, J.J. Harris, C.C. Phillips and R.A. Stradling 184

Magnetotransport properties of Hall-bar with focused-ion-beam written in-plane-gate
R.J. Haug, A.D. Wieck, K. von Klitzing and K. Ploog 192

High-field magneto-resistance in GaAs–GaAlAs heterojunctions
D.R. Leadley, R.J. Nicholas, W. Xu, F.M. Peeters, J.T. Devreese, C.T. Foxon and J.J. Harris 197

Magnetotransport studies of GaSb/InAs crossed gap heterostructures in high magnetic fields
K.S.H. Dalton, R. Bogaerts, R.W. Martin, M. Lakrimi, D.M. Symons, R.J. Warburton, R.J. Nicholas, F. Herlach, N.J. Mason and P.J. Walker 202

Magnetoconductivity probe of inhomogeneity in GaAs–GaAlAs heterojunctions
M. Lakrimi, A.D.C. Grassie, K.M. Hutchings, J.P. Bird, J.J. Harris and C.T. Foxon 206

High-field transport in high carrier density GaAs/$Ga_{0.8}In_{0.2}As$/$Ga_{0.75}Al_{0.25}As$ heterostructures
M. van der Burgt, A. Van Esch, F.M. Peeters, M. Van Hove, G. Borghs and F. Herlach 211

Intersubband scattering in GaAs/$Al_xGa_{1-x}As$ heterostructures
W. de Lange, F.A.P. Blom, P.J. van Hall, P.M. Koenraad and J.H. Wolter 216

Subband population and electron subband mobility for two interacting Si-δ-doping layers in GaAs
P.M. Koenraad, A.C.L. Heessels, F.A.P. Blom, J.A.A.J. Perenboom and J.H. Wolter 221

First evidence of Shubnikov–De Haas oscillations in CdTe
S. Scholl, A. Waag, D. Hommel, K. von Schierstedt, W. Ossau and G. Landwehr 226

Shubnikov–De Haas oscillations in n-$CuInSe_2$
E. Arushanov, L. Essaleh, J. Galibert, J. Leotin and S. Askenazy 229

Experimental study of the energy band structure of $Sc_{1-x}Er_xAs$ layers in pulsed magnetic fields
R. Bogaerts, A. Van Esch, L. Van Bockstal, F. Herlach, F.M. Peeters, F. DeRosa, C.J. Palmstrøm and S.J. Allen Jr. 232

Low-temperature, low-frequency dielectric constant of lightly doped n-InP at high magnetic fields
S. Abboudy 236

Chapter 6. Magneto-tunneling

High magnetic field studies of resonant tunneling via shallow impurities in δ-doped quantum wells
J.W. Sakai, T.M. Fromhold, P.H. Beton, M. Henini, L. Eaves, P.C. Main, F.W. Sheard and G. Hill 241

Electronic perpendicular transport in GaAs–AlAs superlattices in a high magnetic field
F. Aristone, A. Sibille, J.F. Palmier, D.K. Maude, J.C. Portal and F. Mollot 246

Quantum-limit behavior of magneto-tunneling in an AlGaAs/GaAs/AlGaAs double-barrier structure under high magnetic fields up to 40 T
N. Kamata, K. Yamada, N. Miura and L. Eaves 250

High-magnetic-field effects on a resonant-tunneling hot-electron transistor measured up to 40 T
T. Strutz, T. Takamasu, N. Miura and K. Imamura 254

Study of negative differential conductivity in an InAs/GaSb/InAs diode under pulsed high magnetic fields
T. Takamasu, N. Miura, K. Taira, K. Funato and H. Kawai 259

Hamiltonian nonseparability and its consequences in semiconductor heterostructures subjected to high longitudinal magnetic fields
A. Nogaret, L.A. Cury, D.K. Maude, J.C. Portal, D.L. Sivco, A.Y. Cho and G. Hill 263

Chapter 7. Quantum wells and superlattices

High magnetic field studies of the crossed-gap superlattice system InAs/GaSb
R.J. Nicholas, K.S.H. Dalton, M. Lakrimi, C. Lopez, R.W. Martin, N.J. Mason, G.M. Summers, G.M. Sundaram, D.M. Symons, P.J. Walker, R.J. Warburton, M.I. Eremets, D.J. Barnes, N. Miura, L. Van Bockstal, R. Bogaerts and F. Herlach 268

Interacting Landau and Stark quantizations in superlattices under high magnetic fields
A. Alexandrou 277

High magnetic field studies of the confined hole states in quantum wells grown on novel substrate orientations
R.K. Hayden, E.C. Valadares, M. Henini, L. Eaves, D.K. Maude, J.C. Portal and G. Hill 285

Interface optical phonon mode coupling in GaAs/AlAs quantum wells at high magnetic fields
G.Q. Hai, F.M. Peeters and J.T. Devreese 289

Insulator-like miniband transport in heavily doped superlattices under high magnetic fields
H. Noguchi, H. Sakaki, T. Takamasu and N. Miura 293

Chapter 8. Two-dimensional systems

Electronic states and magnetotransport in δ-layer systems
F. Koch 298

Energy level statistics in 2D disordered electron systems subject to strong magnetic fields
H. Kuwano, Y. Ono and K. Slevin 306

Electronic states of a two-dimensional tight-binding model in electric and magnetic fields
T. Nakanishi, T. Ohtsuki and M. Saitoh 310

Negative density of states of a 2DEG in the ultraquantum limit
S.I. Dorozhkin, R.J. Haug, K. von Klitzing and K. Ploog 314

Electronic structure of the triangular quantum well in a tilted magnetic field
S.J. Lee, M.J. Park, G. Ihm, M.L. Falk, S.K. Noh, T.W. Kim and B.D. Choe 318

On negative dielectric permittivity in the quantum Hall regime
H. Nielsen, S.V. Kravchenko, V.M. Pudalov and D.A. Rinberg 323

High-frequency measurements of the magnetoresistance anomaly of kish graphite under pulsed high magnetic fields
T. Takamasu, H. Ochimizu and N. Miura 327

Magnetic-field-induced electronic phase transition in graphite
H. Yaguchi, Y. Iye, T. Takamasu and N. Miura 332

Landau level quantization and superconductivity
H. Akera, A.H. MacDonald and M.R. Norman 337

Chapter 9. Quantum wires

Nonlocal magnetoresistance of diffusive wires in high magnetic fields
P.C. Main, A.K. Geim, P.H. Beton and L. Eaves 341

Conductance fluctuations in a quasi-ballistic narrow wire confined by split metal gates
T. Onishi, Y. Ochiai, M. Kawabe, K. Ishibashi, J.P. Bird, Y. Aoyagi and T. Sugano 351

Analytical study of electrical transport in quantum wires in the presence of magnetic field and spin–orbit scattering
H. Tamura and T. Ando 355

Conductance fluctuations in quantum wires in high magnetic fields
T. Ando 361

Quantum Hall conduction in quantum wires
T. Ando and H. Aoki 365

The observation of Hall plateaux in diffusive quasi-two-dimensional wires of submicron size
M. van der Burgt, A.K. Geim, L. Van Bockstal, S.V. Dubonos and F. Herlach 369

Magnetoconductance fluctuation of a nondegenerate electron gas in silicon
K. Itoh, Y. Sasaki and Y. Iye 374

Chapter 10. Quantum dots

Single-electron capacitance spectroscopy of semiconductor microstructures
R.C. Ashoori, H.L. Störmer, J.S. Weiner, L.N. Pfeiffer, S.J. Pearton, K.W. Baldwin and K.W. West 378

Magic number ground states of quantum dots in a magnetic field
P.A. Maksym 385

Cyclotron resonance of magnetopolarons in anisotropic parabolic quantum dots
R. Haupt and L. Wendler 394

Aharonov–Bohm effect in antidot structures
F. Nihey and K. Nakamura 398

Chapter 11. Shallow impurities

Shallow donors and a fluctuating potential
K. Karpierz, J. Lusakowski, M.L. Sadowski and M. Grynberg 403

Far-infrared magneto-optical studies of D^- ions and many-electron effects on donor impurities in quasi-2D semiconductor structures
J.-P. Cheng, S. Holmes, Y.J. Wang and B.D. McCombe 409

Polaron correction to the D^- center in a quantum well
J.M. Shi, F.M. Peeters and J.T. Devreese 417

Chapter 12. Diluted magnetic semiconductors

Dimensional dependence of antiferromagnetism in diluted magnetic semiconductor heterostructures
W.J. Ossau and B. Kuhn-Heinrich 422

FIR and transport properties of II–VI narrow gap semiconductors in high magnetic fields
M. von Ortenberg 432

An optical study of $Cd_{0.989}Co_{0.011}Se$ in high magnetic fields
M. Inoue, N. Adachi, I. Mogi, G. Kido, Y. Nakagawa and Y. Oka 441

Magnetic quantum oscillation of $Hg_{1-x-y}Cd_xMn_yTe$ in the open-gap range grown by LPE
S. Kuroda, H.-J. Kwon and K. Takita 445

High-field magnetization of $(Cd_{1-x-y}Zn_yMn_x)_3As_2$
H. Bednarski, J.C. Portal, J. Cisowski, H.J.G. Draaisma, T. Goto and W. Lubczyński 451

Magnetic-field- and temperature-dependent exciton delocalisation in a $CdTe/Cd_{1-x}Mn_xTe$ superlattice
M.J. Lawless, R.J. Nicholas, B. Lunn and D.E. Ashenford 455

High-field magneto-optical study of localized excitons in $CdTe/Cd_{1-x}Mn_xTe$ multiple quantum wells
H. Akinaga, Y. Shirai, S. Kuroda, K. Takita, S. Sasaki and N. Miura 460

New aspects of resonant transitions in the mixed-valence compound Hg(Fe)Se
O. Portugall, J. Kono, T. Takamasu, Y. Shimamoto, N. Miura, R. Schoenfeld, M. von Ortenberg and W. Dobrowolski 465

Chapter 13. Organic conductors

Magneto-optical and magneto-transport studies of electron–electron interactions in organic conductors using fields up to 50 T
J. Singleton, F.L. Pratt, M. Doporto, J.M. Caulfield, S.O. Hill, T.J.B.M. Janssen, I. Deckers, G. Pitsi, F. Herlach, W. Hayes, J.A.A.J. Perenboom, M. Kurmoo and P. Day 470

High-field magnetotransport phenomena in organic low-dimensional conductors
T. Osada, S. Kagoshima and N. Miura 481

Fermi surface and magnetic properties of low-dimensional organic conductors
J.S. Brooks, S.J. Klepper, C.C. Agosta, M. Tokumoto, N. Kinoshita, Y. Tanaka, S. Uji, H. Aoki, A.S. Perel, G.J. Athas, X. Chen, D.A. Howe and H. Anzai 489

An anomalous magnetoresistance in α-$(BEDT–TTF)_2I_3$ under high pressure: a possible new phase
T. Ojiro, K. Kajita, Y. Nishio, H. Kobayashi, A. Kobayashi, R. Kato and Y. Iye 494

On the anomalous magnetoresistance in α-$(BEDT–TTF)_2I_3$
Y. Hasegawa and H. Fukuyama 498

Theory of field-induced spin-density wave states in Bechgaard salts: $(TMTSF)_2X$
Y. Hori, K. Machida and M. Nakano 503

Magnetic quantum oscillations in low-dimensional organic conductors (BEDT–TTF)$_2$X
M. Tokumoto, A.G. Swanson, J.S. Brooks, C.C. Agosta, S.T. Hannahs, N. Kinoshita, H. Anzai, M. Tamura, H. Tajima, H. Kuroda, A. Ugawa and K. Yakushi 508

Chapter 14. High magnetic field techniques

High-performance pulsed-field coils for applications in solid state research
F. Herlach, L. Van Bockstal, G. Heremans and L. Li 513

Relaxation measurements of flux pinning by a double pulsed field
M. Motokawa, H. Nojiri, M. Mino and T. Bokui 522

Chapter 15. Closing address

Semiconductors in high magnetic fields – recent trends
G. Landwehr 526

List of contributors 535

Subject index 539